

On the Silov Boundary of a Pseudoconvex Domain in \mathbb{C}^n with $C^{2+\alpha}$ Boundary

SETSUO TANIGUCHI

*Department of Applied Science, Faculty of Engineering,
Kyushu University, Fukuoka 812, Japan*

Communicated by Paul Malliavin

Received March 1990

Let D be a bounded pseudoconvex domain in \mathbb{C}^n and $\partial_{\text{spc}} D$ be the totality of strictly pseudoconvex boundary points. When D has a $C^{2+\alpha}$ plurisubharmonic defining function, a holomorphic diffusion process which never approaches $\partial D \setminus \overline{\partial_{\text{spc}} D}$ is constructed. This diffusion process is used to show that the Silov boundary of D coincides with $\overline{\partial_{\text{spc}} D}$. © 1991 Academic Press, Inc.

1. INTRODUCTION

Let D be a bounded pseudoconvex domain in \mathbb{C}^n . The Silov boundary $S(D)$ of D is the smallest closed subset of the boundary ∂D where the maximum principle holds for all functions h holomorphic on D and continuous up to the boundary: $\sup_{z \in D} |h(z)| = \sup_{z \in S(D)} |h(z)|$. Let $\partial_{\text{spc}} D$ be the totality of strictly pseudoconvex boundary points. For the definition, see the end of Section 2. When D has a C^3 plurisubharmonic (abbreviated psh) defining function, Debiard and Gaveau [1] have shown that $S(D)$ coincides with the closure $\overline{\partial_{\text{spc}} D}$ by constructing a suitable Kähler diffusion process. In this paper, employing another holomorphic diffusion process, we see that $S(D) = \overline{\partial_{\text{spc}} D}$ in the case where D has a $C^{2+\alpha}$ psh defining function for some $0 < \alpha \leq 1$.

A holomorphic diffusion process $\mathbb{M} = (Z_t, \zeta, P_z)$ on D is by definition a $C_0^\infty(D)$ -regular symmetric diffusion process with the lifetime ζ such that $h(Z_t \wedge \tau_K)$ is a P_z -martingale for \mathbb{M} -q.e. $z \in D$, where h is holomorphic on D , K is a compact set in D , $\tau_K = \inf\{t > 0 : Z_t \notin K\}$, and by “ \mathbb{M} -q.e.” we mean “except for a set of zero capacity with respect to the 1-capacity associated with \mathbb{M} .” For the definition of symmetric diffusion processes and the associated 1-capacity, see [2]. A Kähler diffusion process, the minimal diffusion process generated by $\Delta/2$, Δ being the Laplace–Beltrami operator associated with a Kähler metric, is a typical example of holomorphic diffusion processes.

By the martingale convergence theorem, we obtain

$$P_z[Z_{\zeta_-} \equiv \lim_{t \uparrow \zeta} Z_t \text{ exists}] = 1, \quad \mathbb{M}\text{-q.e. } z \in D, \quad (1.1)$$

since D is bounded. Then, for each h holomorphic on D and continuous up to ∂D , it holds that

$$h(z) = E_z[h(Z_{\zeta_-})], \quad \mathbb{M}\text{-q.e. } z \in D,$$

where E_z stands for the expectation with respect to P_z . Suppose that

$$P_z[Z_{\zeta_-} \in \partial D] = 1, \quad \mathbb{M}\text{-q.e.} \quad (1.2)$$

Then, if we define an open subset $\Gamma(\mathbb{M})$ of ∂D by

$$\Gamma(\mathbb{M}) = \bigcup \{U : U \text{ is an open set in } \mathbb{C}^n \text{ such that } U \cap \partial D \neq \emptyset \text{ and } P_z[Z_{\zeta_-} \in U \cap \partial D] = 0, \mathbb{M}\text{-q.e. } z \in D\}, \quad (1.3)$$

then $S(D) \subset D \setminus \Gamma(\mathbb{M})$. Thus, if $\partial D \setminus \overline{\partial_{\text{spc}} D} \subset \Gamma(\mathbb{M})$ then $S(D) = \overline{\partial_{\text{spc}} D}$, because it is well known that $\overline{\partial_{\text{spc}} D} \subset S(D)$ [9]. It is this argument that is taken advantage of by Debiard and Gaveau [1]. Indeed, they have constructed a Kähler diffusion process \mathbb{M}_0 such that $\partial D \setminus \overline{\partial_{\text{spc}} D} \subset \Gamma(\mathbb{M}_0)$ in the case where D has a C^3 psh defining function. Motivated by their work, the author studied the boundary behaviour of holomorphic diffusion processes in [6, 7] and its applications to the complex Monge–Ampère operator. However, it has been open whether there is a holomorphic diffusion process which does not approach $\partial D \setminus \overline{\partial_{\text{spc}} D}$ when the regularity of the boundary is less than C^3 . In the present paper, our aim is to answering the question affirmatively:

THEOREM. *Assume that there exist a domain Ω , a function $\sigma \in C^2(\Omega)$, which is psh on Ω , and a real number $0 < \alpha \leq 1$ such that $\bar{D} \subset \Omega$, $D = \{\sigma < 0\}$, $d\sigma \neq 0$ on ∂D , and $\partial^2 \sigma / \partial z^i \partial \bar{z}^j$, $1 \leq i, j \leq n$, are all α -Hölder continuous. Then, there exists a holomorphic diffusion process \mathbb{M} such that $\partial D \setminus \overline{\partial_{\text{spc}} D} \subset \Gamma(\mathbb{M})$.*

It follows from the theorem that

COROLLARY. *Let D satisfy the same assumption as stated in the theorem. Then $S(D) = \overline{\partial_{\text{spc}} D}$.*

2. PRELIMINARIES

Let G be a bounded domain in \mathbb{C}^n . A C^2 function σ on G is said to be psh if the matrix $((\partial^2 \sigma / \partial z^i \partial \bar{z}^j)(z))_{1 \leq i, j \leq n}$ is nonnegative definite at each

$z \in G$. If the matrix is positive definite at every $z \in G$, σ is called strictly psh. We establish that:

LEMMA 2.1. *Let $\sigma, \tau \in C^2(G)$ be both negative and psh on G . Then, $-(\sigma)^\delta (\tau)^\varepsilon$ is psh for any $\delta > 0$ and $\varepsilon > 0$ with $\delta + \varepsilon \leq 1$. Moreover, $-(\sigma)^\delta (\tau)^\varepsilon$ is strictly psh if either σ or τ is so.*

Proof. For the sake of simplicity, for a C^2 function f , we write $\partial\bar{\partial}f$, and $\partial f, \bar{\partial}f$ for the matrix $(\partial^2 f / \partial z^i \partial \bar{z}^j)_{1 \leq i, j \leq n}$ and the vectors $(\partial f / \partial z^i)_{1 \leq i \leq n}, (\bar{\partial} f / \partial \bar{z}^i)_{1 \leq i \leq n}$, respectively. For $\xi = (\xi^1, \dots, \xi^n), \eta = (\eta^1, \dots, \eta^n) \in \mathbb{C}^n$, $\xi \cdot \eta$ denotes the matrix $(\xi^i \eta^j)_{1 \leq i, j \leq n}$.

Let $p = -(\sigma)^\delta (\tau)^\varepsilon$. Since

$$\partial p = (-p) \left(\frac{\delta}{-\sigma} \partial \sigma + \frac{\varepsilon}{-\tau} \partial \tau \right),$$

by differentiating $-\log(-p)$, we obtain

$$\begin{aligned} \frac{1}{-p} \partial \bar{\partial} p &= \frac{\varepsilon}{-\sigma} \partial \bar{\partial} \sigma + \frac{\varepsilon}{-\tau} \partial \bar{\partial} \tau + \frac{\delta - \delta^2}{(-\sigma)^2} \partial \sigma \cdot \bar{\partial} \sigma + \frac{\varepsilon - \varepsilon^2}{(-\tau)^2} \partial \tau \cdot \bar{\partial} \tau \\ &\quad - \frac{\delta \varepsilon}{\sigma \tau} (\partial \sigma \cdot \bar{\partial} \tau + \partial \tau \cdot \bar{\partial} \sigma). \end{aligned} \quad (2.1)$$

Let us denote by \langle, \rangle the standard inner product in \mathbb{C}^n and set $L_f(z : \xi) = \sum_{i,j=1}^n \xi^i (\partial^2 f / \partial z^i \partial \bar{z}^j)(z) \bar{\xi}^j$ for $f \in C^2(G)$ and $\xi = (\xi^1, \dots, \xi^n) \in \mathbb{C}^n$. Then, (2.1) yields

$$\begin{aligned} \frac{1}{-p} L_p(z : \xi) &\geq \frac{\delta}{-\sigma} L_\sigma(z : \xi) + \frac{\varepsilon}{-\tau} L_\tau(z : \xi) \\ &\quad + (\delta - \delta^2) \left\{ \left| \left\langle \xi, \frac{1}{-\sigma} \bar{\partial} \sigma(z) \right\rangle \right| - \frac{\varepsilon}{1 - \delta} \left| \left\langle \xi, \frac{1}{-\tau} \bar{\partial} \tau(z) \right\rangle \right| \right\}^2 \\ &\quad + \frac{\varepsilon(1 - \delta - \varepsilon)}{1 - \delta} \left| \left\langle \xi, \frac{1}{-\tau} \bar{\partial} \tau(z) \right\rangle \right|^2. \end{aligned}$$

The assertion from this inequality follows immediately.

The identity (2.1) also implies the following:

LEMMA 2.2. *Let $\sigma, \tau \in C^2(D)$ be both negative and psh on G . Suppose that $-1 \leq \sigma$ and that*

$$M \equiv \sup \left\{ |\tau(z)|, \left| \frac{1}{\tau(z)} \right|, |\partial \tau(z)|, \|\partial \bar{\partial} \tau(z)\| : z \in D \right\} < +\infty,$$

where for a \mathbb{C} -valued $n \times n$ -matrix A , $\|A\|^2 = \text{trace } AA^*$. Then, it holds that

$$\begin{aligned} & \text{the minimal eigenvalue of } \partial\bar{\partial}(-(-\sigma)^\delta(-\tau)^\epsilon)(z) \\ & \leq M^\epsilon \{ \delta + \varepsilon(M^2 + M^3) \} (-\sigma)^{\delta-1} \{ L_\sigma(z : \xi) + (-\sigma(z)) \} \end{aligned}$$

for every $z \in G$ and $\xi \in \mathbb{C}^n$ with $|\xi| = 1$ and $\langle \xi, \bar{\partial}\sigma(z) \rangle = 0$.

We end this section by recalling the definition of strictly pseudoconvex points. A domain G is said to possess a C^2 psh defining function if there are an open set O in \mathbb{C}^n and $\psi \in C^2(O)$ such that $\bar{G} \subset O$, $G \cap O = \{\psi < 0\}$ and $d\psi \neq 0$ on ∂G . Such a function ψ is called a C^2 psh defining function of G . For a bounded domain G possessing a C^2 psh defining function ψ , we set

$$\partial_{\text{spc}} G = \{z \in \partial G : L_\psi(z : \xi) > 0 \text{ for every } \xi \in \mathbb{C} \setminus \{0\} \text{ with } \langle \xi, \bar{\partial}\psi(z) \rangle = 0\}.$$

3. PROOF OF THEOREM

We first fix the notation which we will use. Without loss of generality we may assume that

$$-1 \leq \sigma(z) < 0, \quad z \in D. \quad (3.1)$$

Take $0 < \beta, \gamma < 1$, and $R > 0$ such that

$$\alpha + (\beta\gamma - 1)(n-1) > 0 \quad (3.2)$$

$$R > 2 \sup\{|z|^2 : z \in D\}. \quad (3.3)$$

We set

$$p = -(-\sigma)^\beta (R - |z|^2)^{1-\beta}. \quad (3.4)$$

By Lemma 2.1, p and $-(-p)^\gamma$ are both strictly psh on D . Define a positive current θ of bidegree $(n-1, n-1)$ by

$$\theta(dz^i \wedge \sqrt{-1} d\bar{z}^j) = a^{ij} dV, \quad 1 \leq i, j \leq n, \quad (3.5)$$

where (a^{ij}) is the cofactor matrix of $\partial\bar{\partial}(-(-p)^\gamma)$ and V is the Lebesgue measure on D . Let m be an everywhere dense positive Radon measure on D given by

$$dm = (-\sigma)^{\alpha + (\beta\gamma - 1)(n-1) - 2} dV. \quad (3.6)$$

We define a symmetric form \mathcal{E}_θ by

$$\mathcal{E}_\theta(f, g) = \int_D df \wedge d^c g \wedge \theta, \quad f, g \in C_0^\infty(D).$$

Note that

$$\theta = \frac{1}{2^{2n-1}(n-1)!} \{dd^c(-(-p)^\gamma)\}^{n-1} \quad (3.7)$$

and hence θ is closed. It follows from (3.5), (3.6), and the strict plurisubharmonicity of $-(-p)^\gamma$ that for every relatively compact open $G \subset D$,

$$dd^c |z|^2 \wedge \theta \leq C_G m \quad \text{on } G$$

for some $C_G > 0$. Thus, by an argument similar to that in [4, Lemma 1], we see that \mathcal{E}_θ is closable on $L^2(D : m)$. The closure \mathcal{E} is a $C_0^\infty(D)$ -regular, local, Dirichlet form on $L^2(D : m)$ such that every holomorphic function h on D is \mathcal{E} -harmonic. \mathcal{E} then admits a unique holomorphic diffusion process $\mathbb{M} = (Z_t, \zeta, P_z)$ up to equivalence. See [4, 8]. We now show

LEMMA 3.1. \mathbb{M} explodes:

$$P_z[\zeta < +\infty] = 1, \quad \mathbb{M}\text{-}q.e. \ z \in D.$$

In particular, (1.2) is satisfied.

Proof. Let $(\mathcal{F}, \mathcal{E})$ be the Dirichlet space of \mathbb{M} and $(\mathcal{F}', \mathcal{E}')$ be that of the absorbing barrier Brownian motion on D . Then, for every relatively compact open set G in D , there is a constant $C_0 > 0$ such that

$$\mathcal{E}(f, f) \geq C_0 \mathcal{E}'(f, f), \quad f \in C_0^\infty(G)$$

and

$$C_0 V \leq m \leq C_0^{-1} V.$$

By [3, Proposition 1.5], we see that \mathbb{M} is irreducible. Then, as is seen in [7, 10], \mathbb{M} is transient.

We now recall that if $D_0 = \{\varphi < 0\}$ is a bounded pseudoconvex domain in \mathbb{C}^n with a continuous psh defining function φ and \mathbb{N} is a transient holomorphic diffusion process with the symmetrising measure m_0 , then \mathbb{N} explodes whenever $\int_{D_0} |\varphi| dm_0 < \infty$. See [10, Theorem 1.1]. Because of (3.2), it is elementary to see that

$$\int_D |\sigma| dm < +\infty.$$

Thus, we obtain that \mathbb{M} explodes.

To see that $\partial D \setminus \overline{\partial_{\text{spc}} D} \subset \Gamma(\mathbb{M})$, we fix an arbitrary $z^* \in \partial D \setminus \overline{\partial_{\text{spc}} D}$ and an open set U such that $U \cap \partial D \subset \partial D \setminus \overline{\partial_{\text{spc}} D}$. Shrinking U if necessary, we may assume that a mapping $\Phi : (\partial D \cap U) \times (-\delta_0, \delta_0) \rightarrow U$ given by

$$\Phi(w, a) = w + an_w,$$

n_w being the outer normal vector at w , is a C^2 -diffeomorphism for some $\delta_0 > 0$. For $z \in U$, $w \in \partial D$ satisfying $\Phi(w, a) = z$ is denoted by $\pi(z)$. Shrinking U again if necessary, we may, moreover, assume that there are $C_1, C_2 > 0$ such that $C_1 \delta \leq \frac{1}{2}$ and

$$\partial\sigma(z) \neq 0 \quad (3.8)$$

$$|\partial\sigma(z)|^{-1} |\delta\sigma(z) - 4\langle \delta\sigma(\pi(z)), \delta\sigma(z) \rangle \delta\sigma(\pi(z))| \leq C_1 |z - \pi(z)| \quad (3.9)$$

$$|z - \pi(z)| \leq C_2 |\sigma(z)| \quad (3.10)$$

for every $z \in U$. On such a U , we obtain the following inequality, which is verified by modifying the argument used in [1].

LEMMA 3.2. *Let U be as above. Then, for some $C_3 > 0$, it holds that*

$$C_3 dd^c |z|^2 \wedge \theta \geq (-\sigma)^{-\beta\gamma - \alpha + 1} dd^c (-(-p)^\gamma) \wedge \theta \quad \text{on } U \cap D. \quad (3.11)$$

Proof. Take an arbitrary $z \in U \cap D$ and fix it. Choose $\xi \in \mathbb{C}^n$ such that $|\xi| = 1$, $\langle \xi, \delta\sigma(\pi(z)) \rangle = 0$, and $L_\sigma(\pi(z) : \xi) = 0$. We put

$$\eta = \xi - |\delta\sigma(z)|^{-2} \langle \xi, \delta\sigma(z) \rangle \delta\sigma(z).$$

Since $\langle \xi, \delta\sigma(\pi(z)) \rangle = 0$ and $|\delta\sigma(\pi(z))| = \frac{1}{2}$, it follows from (3.9) that

$$\begin{aligned} |\eta - \xi| &\leq |\delta\sigma(z)|^{-1} |\delta\sigma(z) - 4\langle \delta\sigma(z), \delta\sigma(\pi(z)) \rangle \delta\sigma(\pi(z))| \\ &\leq C_1 |z - \pi(z)| \\ &\leq C_1 \delta \leq \frac{1}{2}. \end{aligned} \quad (3.12)$$

In particular,

$$\frac{1}{2} \leq |\eta| \leq \frac{3}{2}. \quad (3.13)$$

Define constants C_4 and C_5 by

$$\begin{aligned} C_4 &= \sup_{u, v \in U} \frac{\|\partial\delta\sigma(u) - \partial\delta\sigma(v)\|}{|u - v|^\alpha} < \infty \\ C_5 &= \sup_{w \in U} \|\partial\delta\sigma(w)\| < \infty. \end{aligned}$$

Then, by a straightforward computation, we obtain that

$$\begin{aligned} L_\sigma(z : \eta) &= L_\sigma(z : \eta) - L_\sigma(\pi(z) : \xi) \\ &\leq (|\eta| + 1) C_5 |\eta - \xi| + C_4 |z - \pi(z)|^\alpha. \end{aligned}$$

Combining this with (3.10), (3.12), and (3.13), we have

$$L_\sigma\left(z : \frac{\eta}{|\eta|}\right) \leq 2(5C_1 C_5 + 2C_4) C_2 |\sigma(z)|^\alpha. \quad (3.14)$$

Because of Lemma 2.2, (3.14) implies the existence of a constant $C_6 > 0$, independent of z , such that

$$\text{the minimal eigenvalue of } \partial\bar{\partial}(-(-p)^\gamma)(z) \leq C_6(-\sigma(z))^{\beta\gamma + \alpha - 1}. \quad (3.15)$$

It follows from (3.5) and (3.7) that

$$dd^c |z| \wedge \theta = 2 \operatorname{trace}(a^{i\bar{j}}) dV, \quad (3.16)$$

$$dd^c(-(-p)^\gamma) \wedge \theta = 2n \det(\partial\bar{\partial}(-(-p)^\gamma)) dV. \quad (3.17)$$

If we set $(b^{i\bar{j}}) = (\partial\bar{\partial}(-(-p)^\gamma))^{-1}$, then (3.15) implies that

$$\begin{aligned} \operatorname{trace}(a^{i\bar{j}}) &= \operatorname{trace}(b^{i\bar{j}}) \det(\partial\bar{\partial}(-(-p)^\gamma)) \\ &\geq C_6^{-1} (-\sigma)^{-\beta\gamma - \alpha + 1} \det(\partial\bar{\partial}(-(-p)^\gamma)) \quad \text{on } U \cap D. \end{aligned}$$

Combining this with (3.16), (3.17), we obtain the desired inequality.

We now consider the relationship between the measures m and $dd^c(-\log(-p)) \wedge \theta$. We verify

LEMMA 3.3. *Let U be as before. Then, there is a constant C_7 such that*

$$dd^c(-\log(-p)) \wedge \theta \leq C_7 m \quad \text{on } U \cap D. \quad (3.18)$$

Proof. It is easy to show that

$$dd^c(-\log(-p)) \wedge \theta \leq \gamma(1 - \gamma) R^{-\beta\gamma} (-\sigma)^{-\beta\gamma} dd^c(-(-p)^\gamma) \wedge \theta \quad \text{on } D. \quad (3.19)$$

Plugging (3.11) into (3.19), we obtain that

$$dd^c(-\log(-p)) \wedge \theta \leq \gamma(1 - \gamma) R^{-\beta\gamma} C_3 (-\sigma)^{\alpha - 1} dd^c |z|^2 \wedge \theta \quad \text{on } U \cap D. \quad (3.20)$$

Take an arbitrary $z \in U \cap D$ and fix it. Choose $\xi_1, \dots, \xi_n \in \mathbb{C}^n$ such that $\langle \xi_i, \xi_j \rangle = \delta_{ij}$ and $\langle \xi_j, \delta\sigma(z) \rangle = 0$, $1 \leq j \leq n-1$. It follows from (2.1) that there exists a constant C_8 , independent of z , such that

$$|\langle \xi_i, \delta\hat{\partial}(-(-p)^\gamma)(z) \xi_j \rangle| \leq C_8(-\sigma(z))^{\beta\gamma-1} \quad \text{if } i+j < 2n \quad (3.21)$$

and

$$|\langle \xi_n, \delta\hat{\partial}(-(-p)^\gamma)(z) \xi_n \rangle| \leq C_8(-\sigma(z))^{\beta\gamma-2}. \quad (3.22)$$

Let (c^{ij}) be the cofactor matrix of $(\langle \xi_i, \delta\hat{\partial}(-(-p)^\gamma)(z) \xi_j \rangle)_{1 \leq i, j \leq n}$. Note that

$$\text{trace}(a^{ij}) = \text{trace}(c^{ij}).$$

Hence, (3.16), (3.21), and (3.22) imply that

$$dd^c |z|^2 \wedge \theta \leq C_9(-\sigma)^{(\beta\gamma-1)(n-1)-1} dV \quad \text{on } U \cap D$$

for some constant C_9 . Combining this with (3.20), we obtain that (3.18) holds.

We finally investigate the behaviour of \mathbb{M} on $U \cap D$ in the following lemma. The assertion of the theorem is an immediate consequence of the lemma, because of the maximality of $I(\mathbb{M})$.

LEMMA 3.4. *Let U be as above. Then, it holds that*

$$P_z[Z_{\zeta-} \in \partial M \cap U] = 0 \quad \mathbb{M}\text{-q.e. } z \in D.$$

Proof. Recall that $-\log(-p)$ is locally in \mathcal{F} , where \mathcal{F} is the domain of the Dirichlet form associated with \mathbb{M} . See [4, 5]. The continuous additive functional (abbreviated CAF) $-\log(-p(Z_t)) + \log(-p(Z_0))$ is decomposed as

$$-\log(-p(Z_t)) + \log(-p(Z_0)) = M_t + N_t, \quad t < \zeta,$$

where M_t is a local martingale CAF and N_t is a CAF of local energy zero (cf. [2]). By the standard time change argument, we have that

$$-\log(-p(Z_t)) + \log(-p(Z_0)) = B(\langle M_t \rangle) + N_t, \quad t < \zeta, \quad (3.23)$$

where $\langle M \rangle$ is the quadratic variation process of M and $B(t)$ is an \mathbb{R}^1 -valued Brownian motion with $B(0) = 0$.

The Revuz measures of N and $\langle M \rangle$ are $dd^c(-\log(-p)) \wedge \theta$ and $d(-\log(-p)) \wedge d^c(-\log(-p)) \wedge \theta$, respectively. See [4, 6]. Since the latter measure dominates the former one, it follows from (3.18) that

$$\langle M \rangle_t \leq N_t \leq C_7 t, \quad t < \zeta \wedge t', \quad (3.24)$$

under P_z , \mathbb{M} -q.e. $z \in U \cap D$, where $\tau' = \inf\{t > 0 : Z_t \in D \setminus U\}$. Equation (3.24) and Lemma 3.1 yield that

$$P_z[\lim_{t \uparrow \zeta \wedge \tau'} \langle M \rangle_t < \infty, \lim_{t \uparrow \zeta \wedge \tau'} N_t < \infty] = 1, \quad \mathbb{M}\text{-q.e. } z \in U \cap D.$$

Combined with (3.23), this implies that

$$P_z[Z_t \in D \cap U, t < \zeta] = 0, \quad \mathbb{M}\text{-q.e. } z \in U \cap D.$$

Hence there exists a Borel set $E \subset D$ such that E is of zero capacity with respect to the 1-capacity associated with \mathbb{M} ,

$$P_z[Z_t \in D \setminus E] = 1, \quad z \in D \setminus E, \quad (3.25)$$

$$P_z[Z_t \in D \cap U, t < \zeta] = 0, \quad z \in U \setminus E. \quad (3.26)$$

Set

$$A_r = \{\zeta > r \text{ and } Z_t \in D \cap U \text{ for } r \leq t < \zeta\}.$$

By the Markov property, (3.25), and (3.26), we obtain that

$$P_z[A_r] = 0 \quad \text{for } z \in D \setminus E,$$

Since $\{Z_{\zeta_-} \in \partial D \cap U\} \subset \bigcup_r A_r$, where the union is taken over all non-negative rational numbers, this implies that

$$P_z[Z_{\zeta_-} \in \partial D \cap U] = 0, \quad \mathbb{M}\text{-q.e. } z \in D.$$

REFERENCES

1. A. DEBIARD AND B. GAVEAU, Frontière de Silov de domaines faiblement pseudoconvexes de C^n , *Bull. Sci. Math.* **100** (1976), 17–31.
2. M. FUKUSHIMA, “Dirichlet Forms and Markov Processes,” Kodansha/North-Holland, Tokyo/Amsterdam, 1980.
3. M. FUKUSHIMA, On recurrence criteria in the Dirichlet space theory, in “From Local Times to Global Geometry, Control and Physics” (K. D. Elworthy, Ed.), Pitman Res. Notes Math. Ser. 150, Longman, Essex, 1987.
4. M. FUKUSHIMA AND M. OKADA, On conformal martingale diffusions and pluripolar sets, *J. Funct. Anal.* **55** (1984), 377–388.
5. M. FUKUSHIMA AND M. OKDADA, On Dirichlet forms for plurisubharmonic functions, *Acta Math.* **159** (1987), 171–213.
6. H. KANEKO AND S. TANIGUCHI, A stochastic approach to the Silov boundary, *J. Funct. Anal.* **74** (1987), 415–429.

7. H. KANEKO AND S. TANIGUCHI, Conformal martingale diffusions and Shilov boundaries, in "Potential Theory and Its Related Fields," RIMS Kokyuroku 610, RIMS Kyoto University, Kyoto, 1987.
8. M. OKADA, Espace de Dirichlet généraux en analyse complexe, *J. Funct. Anal.* **46** (1982), 396–410.
9. H. ROSSI, Holomorphically convex sets, *Ann. Math.* **74** (1961), 470–493.
10. S. TANIGUCHI, The explosion property of holomorphic diffusion processes on a bounded pseudoconvex domain in C^n and its applications, *J. Math. Kyoto Univ.*, to appear.